

1-21-2019

CCNE1 amplification is associated with poor prognosis in patients with triple negative breast cancer.

Zi-Ming Zhao

The Jackson Laboratory, ziming.zhao@jax.org

Susan E Yost


Katherine E Hutchinson

Sierra Min Li

Yate-Ching Yuan

See next page for additional authors

Follow this and additional works at: <https://mouseion.jax.org/stfb2019>

 Part of the [Life Sciences Commons](#), and the [Medicine and Health Sciences Commons](#)

Recommended Citation

Zhao, Zi-Ming; Yost, Susan E; Hutchinson, Katherine E; Li, Sierra Min; Yuan, Yate-Ching; Noorbakhsh, Javad; Liu, Zheng; Warden, Charles; Johnson, Radia M; Wu, Xiwei; Chuang, Jeffrey H; and Yuan, Yuan, "CCNE1 amplification is associated with poor prognosis in patients with triple negative breast cancer." (2019). *Faculty Research 2019*. 28.

<https://mouseion.jax.org/stfb2019/28>

This Article is brought to you for free and open access by the Faculty Research at The Mouseion at the JAXlibrary. It has been accepted for inclusion in Faculty Research 2019 by an authorized administrator of The Mouseion at the JAXlibrary. For more information, please contact ann.jordan@jax.org.

Authors

Zi-Ming Zhao, Susan E Yost, Katherine E Hutchinson, Sierra Min Li, Yate-Ching Yuan, Javad Noorbakhsh, Zheng Liu, Charles Warden, Radia M Johnson, Xiwei Wu, Jeffrey H Chuang, and Yuan Yuan

RESEARCH ARTICLE

Open Access



CCNE1 amplification is associated with poor prognosis in patients with triple negative breast cancer

Zi-Ming Zhao¹, Susan E. Yost², Katherine E. Hutchinson³, Sierra Min Li², Yate-Ching Yuan², Javad Noorbakhsh¹, Zheng Liu², Charles Warden², Radia M. Johnson³, Xiwei Wu², Jeffrey H. Chuang^{1†} and Yuan Yuan^{2**}

Abstract

Background: Triple negative breast cancer (TNBC) is aggressive with limited treatment options upon recurrence. Molecular discordance between primary and metastatic TNBC has been observed, but the degree of biological heterogeneity has not been fully explored. Furthermore, genomic evolution through treatment is poorly understood. In this study, we aim to characterize the genomic changes between paired primary and metastatic TNBCs through transcriptomic and genomic profiling, and to identify genomic alterations which may contribute to chemotherapy resistance.

Methods: Genomic alterations and mRNA expression of 10 paired primary and metastatic TNBCs were determined through targeted sequencing, microarray analysis, and RNA sequencing. Commonly mutated genes, as well as differentially expressed and co-expressed genes were identified. We further explored the clinical relevance of differentially expressed genes between primary and metastatic tumors to patient survival using large public datasets.

Results: Through gene expression profiling, we observed a shift in TNBC subtype classifications between primary and metastatic TNBCs. A panel of eight cancer driver genes (*CCNE1*, *TPX2*, *ELF3*, *FANCL*, *JAK2*, *GSK3B*, *CEP76*, and *SYK*) were differentially expressed in recurrent TNBCs, and were also overexpressed in TCGA and METABRIC. *CCNE1* and *TPX2* were co-overexpressed in TNBCs. DNA mutation profiling showed that multiple mutations occurred in genes comprising a number of potentially targetable pathways including PI3K/AKT/mTOR, RAS/MAPK, cell cycle, and growth factor receptor signaling, reaffirming the wide heterogeneity of mechanisms driving TNBC. *CCNE1* amplification was associated with poor overall survival in patients with metastatic TNBC.

Conclusions: *CCNE1* amplification may confer resistance to chemotherapy and is associated with poor overall survival in TNBC.

Keywords: Triple negative breast cancer, CCNE1, Amplification

Background

Triple negative breast cancer (TNBC) is an aggressive type of breast cancer (BC) characterized by a high rate of recurrence and poor overall survival upon cancer metastasis [1, 2]. Hormone-receptor positive (HR+) BCs and HER2-positive (HER2+) BCs harbor well-defined

and targetable biomarkers. Despite the identification of at least four molecular TNBC subtypes, including basal-like 1 (BL1), basal-like 2 (BL2), mesenchymal (M), and luminal androgen receptor (LAR) [3–5], these subtypes have not proven clinically useful due to the intrinsically complex and heterogeneous biology of TNBCs. With the exception of PARP inhibitors for patients with *BRCA1/2* germline mutations, and unlike HR+ or HER2+ BCs, there is no effective targeted therapy available to treat TNBCs.

* Correspondence: yuyuan@coh.org

[†]Jeffrey H. Chuang and Yuan Yuan contributed equally to this work.

²City of Hope Comprehensive Cancer Center and Beckman Research Institute, 1500 E. Duarte Road, Duarte, CA 91010, USA

Full list of author information is available at the end of the article



© The Author(s). 2019 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

In addition to tumor heterogeneity, the biology of TNBC is further complicated by tumor evolution through selective pressure. The molecular evolution of TNBC as the result of chemotherapy and/or radiation-induced selection pressure is well recognized but poorly understood [6]. Molecular discordance between primary and metastatic TNBCs has been observed, but the degree of biological heterogeneity has not been fully explored [7].

Recurrent/resistant BC may differ from primary tumors at multiple levels. Several previous studies demonstrated phenotypic discordances between primary and metastatic tumors in the standard-of-care biomarkers estrogen receptor (ER), progesterone receptor (PR), and HER2 [8, 9]. Discordance of other molecular markers such as *PIK3CA* mutations was also identified [9, 10]. Cyclin E1 (*CCNE1*), along with its catalytic subunit CDK2, plays a critical role in cell cycle regulation, DNA replication, chromosome segregation, and the G1 to S-phase transition [11, 12]. *CCNE1* amplification is associated with primary treatment resistance in high-grade serous ovarian carcinomas (HGSCs) and co-amplification of *TPX2* with *CCNE1* was common [13, 14]. *CCNE1* amplification is also associated with resistance to HER2-targeted therapy in HER2+ BC [15]. However, the role of *CCNE1* in TNBC is not well understood.

Genomic and transcriptomic discordance between primary and metastatic tumors have the potential to reveal novel drivers of metastatic progression, and aid in the selection of late-line therapies [16]. Genomic sequencing has revealed subclonal diversity of primary BCs and chemotherapy-resistant BCs in experimental models [17], but such findings have not been shown in patient tumor samples. Available large-scale public genomic databases from The Cancer Genome Atlas (TCGA) project [18] and the Molecular Taxonomy of Breast Cancer International Consortium (METABRIC) [19] have significantly improved our understanding of breast cancer tumor biology. These data sets contain only newly diagnosed and chemotherapy naïve tumors as opposed to complementary longitudinal primary and metastatic samples. Therefore, genomic characterization of longitudinal samples has the potential to identify novel driver mutations and therapeutic targets.

The primary goal of this study is to characterize the genomic changes between paired primary and metastatic TNBCs through sequencing and mRNA expression profiling, and to identify genomic alterations which may contribute to chemotherapy resistance.

Methods

Cohort of paired primary and recurrent/metastatic TNBC specimens

Paired primary and recurrent TNBC specimens were identified through an IRB-approved protocol from patients with

recurrence between 2002 and 2015. Eligible patients had the following features: stage I - III TNBC; at least one tumor biospecimen available from initial surgery or biopsy; at least one specimen available from a recurrent/metastatic disease; and treatment and clinical outcome data available for chart review. Two cohorts of patients were studied: "Pilot-TNBC" ($n = 10$) and "Discovery-TNBC" ($n = 55$). All pathology samples were formalin-fixed and paraffin-embedded (FFPE). Demographic data such as age, gender, date of birth, date of diagnosis, date of relapse, and date of death or last follow-up (if applicable) were obtained for *CCNE1* amplified patients ($n = 13$, 6 from "Discovery-TNBC", and 7 from clinical report). Disease characteristics such as tumor grade, TNM stage, and ER/PR/HER2 status, as well as treatment variables including surgery, chemotherapy, and radiation therapy were also obtained.

Gene expression profiling of paired TNBCs

Paired primary and recurrent TNBC samples were profiled using the Affymetrix Human GeneChip® 2.0 in the "Pilot-TNBC" cohort. Robust multi-array averages (RMA [20]) were calculated from mRNA expression, summarized at the gene symbol level (23,945 gene symbols) based upon probe-set annotations (instead of transcript clusters) with the Affymetrix Power Tools software [21]. Messenger RNA (mRNA) expression across control probe transcript clusters was also summarized (27,293 control probes). Gene symbol annotations for probe sets were downloaded from the Affymetrix website. When comparing expression between primary and recurrent samples, p -values were calculated using the limma R package [22] and false discovery rate (FDR) values were calculated using the method of Benjamini and Hochberg [23]. The number of differentially expressed genes was maximized by considering genes with $\text{RMA} > 2$ in at least 50% of samples, $|\text{fold-change}| > 1.5$ between averaged recurrent and primary tumors, and $\text{FDR} < 0.25$. The ratio of gene expression between paired recurrent and primary TNBCs was reported as fold-change. Heat maps and clustering of differentially expressed genes were created using the 'seaborn' package under Jupyter Notebook Python 3. For better visualization, we only reported fold changes between -10 and 10 . TNBC subtyping was performed using the online TNBC Type tool from Vanderbilt University Medical Center [24]. Of the "Discovery-TNBC" cohort ($n = 55$), RNA isolation and expression profiling were performed for 35 paired primary-metastatic specimens. Total RNA was isolated from FFPE specimens using the QIAGEN miRNeasy FFPE kit. RNA integrity (RIN) and DV200 scores were determined using an Agilent Bioanalyzer. Per Illumina protocol, TruSeq RNA Access Library Preparation was performed, and libraries were enriched for mRNA

fraction by positive selection, prior to paired-end 2×100 sequencing on the Illumina HiSeq. Using the R package “edgeR version 3.20.9” [25, 26], raw transcript counts were converted to log2-counts per million (log2CPM), dataset-wide lowly- or non-expressed genes were removed, and transcript expression was normalized to ensure similar expression distributions across the dataset.

Targeted exome sequencing of TNBCs

Genomic alterations in FFPE specimens from primary and recurrent TNBCs were detected using the FoundationOne™ sequencing panel. FoundationOne™ identifies base substitutions, insertions and deletions (indels), amplifications with copy number ≥ 6 , and rearrangements. The FoundationOne™ sequencing panel version used herein included the entire coding regions of 395 cancer-related genes, and select introns of 31 genes that are rearranged or altered in cancer, capable of achieving a median sequencing depth greater than 500X [27, 28].

Comparison of gene expression using public databases

Gene expression of 1904 samples from METABRIC (299 TNBC and 1605 non-TNBC) using Illumina HT-12 arrays were downloaded [19]. Whole-transcriptome sequences of 1091 primary breast cancer (115 TNBC and 976 non-TNBC) and 112 normal cases in TCGA were downloaded in the format of ‘illuminahtseq_rnaseqv2-RSEM_genes_normalized (MD5)’ from FireBrowse [18]. Boxplots for gene expression comparisons in METABRIC [19] and TCGA [18] datasets were created using the R “ggplot” package. TCGA data were transformed to log2 of the gene expression value plus 1, to make them the same scale as the METABRIC gene expression data. *P*-values were independently calculated using Wilcoxon and

ANOVA tests to compare between two, and among multiple groups, respectively.

Statistical analysis

Kaplan-Meier (K-M) curves were generated for overall survival (OS) and relapse free survival (RFS), where OS is defined as time from surgery to death, and RFS is defined as time from surgery to disease recurrence. The log-rank test was used to examine survival (OS or RFS) difference based on *CCNE1* copy number alteration ($CN \geq 6$ vs. $CN < 6$), or *CCNE1* mRNA expression (median gene expression as a bifurcation).

Results

Patient characteristics and treatment history

The clinical characteristics, pathological features, treatment histories, and survival of the “Pilot-TNBC” cohort are described in Table 1. The majority of the tumors were infiltrating ductal carcinomas (IDC) (80%, 8/10), stage I-II (70%, 7/10), and the patients received standard-of-care chemotherapy with anthracycline and/or a taxane-containing regimen. RFS ranged from 2 to 39 months, and overall survival ranged from 9 to 113 months. Molecular subtypes according to the Lehmann/Pietenpol classification were determined: BL-1 ($n = 4$), BL-2 ($n = 1$), M ($n = 3$), and LAR ($n = 2$). Four of ten TNBC pairs displayed a shift in the molecular subtype between the primary and metastatic specimens.


Genomic and transcriptomic profiling of paired TNBCs in the “Pilot-TNBC” cohort

In “Pilot-TNBC,” FoundationOne™ sequencing was successful in 7 paired specimens ($n = 14$) (Fig. 1a). A total of 339 genomic alterations including 73 known

Table 1 Patient characteristics in the “Pilot-TNBC” cohort

Pt ID	Spec# 1	Spec# 2	Histology	Age range	Stage	Lehmann subtypes	(Neo)adjuvant chemo	Radiation	RFS (months)	OS (months)
1	Breast	Lung Met	IDC	60–69	II	BL2 → BL2	AC-T	Yes	34	113
2	Breast	Bone Met	IDC	50–59	II	LAR → LAR	AC	No	17	60
3	Breast	LN Met	IDC	40–49	II	BL1 → BL1	AC	No	39	92
4	LN	Liver Met	IDC	40–49	III	BL1 → LAR	Carbo/Taxol	Yes	10	20
5	Breast	Skin Met	IDC	30–39	III	M → M	AC	Unknown	2	9
6	Brain	Soft tissue Met	IDC	50–59	II	M → M	TAC	No	31	67
7	Breast	LN Met	IDC	50–59	I	BL1 → BL2	Declined	No	26	46
8	Endometrium	LN Met	IDC	30–39	III	M → LAR	TC	Unknown	11	58
9	Breast	Contralateral breast Met	ILC	40–49	II	LAR → BL2	AC-T	Yes	8	23
10	Breast	Brain Met	Metaplastic	50–59	II	BL1 → BL1	AC-T	Unknown	16	58

IDC invasive ductal carcinoma, LN lymph node, RFS relapse-free survival (surgery to first relapse), OS overall survival (surgery to death), ILC invasive lobular carcinoma, AC adriamycin/cyclophosphamide, AC-T adriamycin/cyclophosphamide -> paclitaxel, Carbo/taxol carboplatin/paclitaxel, TAC docetaxel/adriamycin/cyclophosphamide, TC docetaxel /cyclophosphamide, Met metastatic tumor, BL1 basal-like 1, BL2 basal-like 2, M mesenchymal, LAR luminal androgen receptor, Spec specimen




mutation/amplification and 266 variants of unknown significance (VUS) were identified. Genomic alterations were identified in the following signaling pathways: cell cycle, p53, PI3K/mTOR, RAS/MAPK, and RTK/GF (Fig. 1b) [29]. There were no *CCNE1* mutations or amplifications detected in the “Pilot-TNBC” cohort. Detailed genomic alteration data is listed in Additional file 1: Table S3. In addition, there was significant inter-patient genomic heterogeneity but little intra-patient variability. These findings not only confirm the genomic heterogeneity of TNBCs, but also highlight the genomic stability of these tumors over the course of time.

To identify genes driving cancer recurrence, gene expression patterns were studied by comparing the paired TNBCs in “Pilot-TNBC.” We found 455 differentially expressed genes (DEG) between metastatic/primary TNBCs (> 1.5 fold changes), with 317 genes upregulated and 138 genes down-regulated in the metastatic TNBCs compared to their matched primaries (Fig. 2a). From the 455 DEGs, the expression of 8 cancer-related genes (*CCNE1*, *TPX2*, *GSK3B*, *CEP76*, *SYK*, *JAK2*, *ELF3* and *FANCL*) was significantly upregulated in recurrent/metastatic TNBCs compared with matched primaries (Fig. 2b). These genes represent potential cancer driver genes, selected based on putative cancer driver gene lists from literature [27, 30–34] and/or inclusion in

the FoundationOne™ targeted sequencing panel. There was no clear association between expression pattern of these genes and Lehmann/Pietenpol subtypes of TNBC (data not shown).

CCNE1 and TPX2 mRNA expression in primary breast cancer

We next analyzed whether genes upregulated in TNBC metastases had systematic expression patterns across breast cancer datasets. To further understand genomic and transcriptomic expression of *CCNE1* and *TPX2* in primary TNBCs, public databases including METABRIC and TCGA were used. TNBCs exhibited higher mRNA expression of both genes compared with non-TNBCs. In METABRIC, the percentage of tumors with *CCNE1* mRNA overexpression (above median) is significantly increased in TNBCs (42.1%, 126/299) compared with non-TNBCs (3.2%, 51/1605) ($p < 0.0001$). In TCGA, similar results were found in TNBCs (48.7%, 56/115) compared with non-TNBCs (7.2%, 70/976) ($p < 0.0001$). In METABRIC, TNBCs also exhibited higher *TPX2* mRNA overexpression (22.4%, 67/299) compared with non-TNBC (2.4%, 38/1605) ($p < 0.001$). In TCGA, similar results were found for *TPX2* mRNA overexpression in TNBCs (40.9%, 47/115) compared with non-TNBCs (7.6%, 74/976) ($p < 0.0001$) (Fig. 3a and b). In addition, *CCNE1* and *TPX2* are co-overexpressed in TNBC



(METABRIC, $n = 299$, $p \leq 0.001$; TCGA, $n = 115$, $p \leq 0.001$) (Fig. 3c and d). Furthermore, *CCNE1* is significantly co-overexpressed with *TPX2* in “Pilot-TNBC” and “Discovery-TNBC” cohorts ($p < 0.001$) (Additional file 2: Figure S2). However, *CCNE1* or *TPX* mRNA overexpression in TNBCs did not confer a difference in OS (data not shown).

In METABRIC, TNBCs also exhibited significantly higher mRNA expression of *FANCL* ($p < 0.0001$), *SYK* ($p < 0.0001$), *ELF3* ($p < 0.01$), *JAK2* ($p < 0.0001$), and *GSK3B* ($p < 0.0001$) compared with non-TNBCs (Additional file 3: Figure S1a). There was no significant change in *CEP76*. In TCGA, TNBCs exhibited significantly higher mRNA expression of *FANCL* ($p < 0.0001$), *SYK* ($p < 0.0001$), and *CEP76* ($p < 0.0001$) compared with non-TNBCs (Additional file 3: Figure S1b). No significant changes in *ELF*, *JAK2* and *GSK3B* were observed. Further, compared with normal tissue, median mRNA expression of these eight genes was significantly elevated in breast cancer tissue (Additional file 3: Figure S1c).

***CCNE1* amplification is associated with poor overall survival in the “Discovery-TNBC” cohort**

In the “Pilot-TNBC” cohort, *CCNE1* amplification was not observed. Further analysis of an independent cohort of “Discovery-TNBC” revealed *CCNE1* amplification ($CN \geq 6$) in 6 of 55 (10.9%) patients (primary and metastatic tumors). The clinical characteristics and outcomes of these patients with *CCNE1*-amplified TNBC are summarized in Table 2. Patients with *CCNE1*-amplified tumors presented with advanced stage disease and had poor pathological responses to neoadjuvant chemotherapy. None of

these patients had *BRCA1/2* mutations. There is a statistically significant association between *CCNE1* amplification and poor OS ($p = 0.023$) (Fig. 4a), but not RFS ($p = 0.25$) (Fig. 4b).

Of the “Discovery-TNBC” cohort ($n = 55$), whole-transcriptome sequencing (RNAseq) was successful in 35 paired primary-metastatic specimens. Of those, 25 specimens exhibited an increase in *CCNE1* mRNA expression in metastatic samples compared to their primary counterparts. No association between *CCNE1* mRNA over-expression and OS was observed in the “Discovery-TNBC” cohort ($p = 0.45$) (Fig. 4c).

We further examined the association of *CCNE1* gene expression and *CCNE1* gene amplification in both primary and metastatic tumors (Fig. 4d). In tumors with no *CCNE1* amplification ($CN < 6$), the mean and variance of *CCNE1* mRNA expression were comparable between primary and metastatic tumors. However, the mean *CCNE1* expression in metastatic tumors was increased compared to primary tumors (expression = 5.29 vs. 3.87, $SD = 0.49$ vs. 2.02).

CCNE1 amplification was more frequently detected in TNBCs (9%, 27/299 in METABRIC and 10%, 11/111 in TCGA) compared to non-TNBCs (1.7%, 27/1605 in METABRIC, $p < 0.0001$ and 3.5%, 33/962 in TCGA, $p = 0.0055$), but these distinctions did not translate into significant OS differences, likely due to limited sample sizes. In TCGA, *TPX2* amplification was more frequent in TNBCs (8.1%, 9/111) compared to non-TNBCs (1.5%, 14/962) ($p = 0.0004$), but not in METABRIC. Interestingly, reflective of our “Discovery-TNBC” cohort data, *CCNE1* amplification





Fig. 3 TNBCs exhibited higher mRNA expression of *CCNE1* and *TPX2* compared with non-TNBCs (Wilcoxon signed rank test) in (a) METABRIC: TNBC ($n = 299$) and non-TNBC ($n = 1605$) ($p < 0.0001$ for both *CCNE1* and *TPX2*) and in (b) TCGA: TNBC ($n = 115$) and non-TNBC ($n = 976$) ($p < 0.0001$ for both *CCNE1* and *TPX2*), y-axis is the log2 of TPM values by RSEM. In TNBCs, *CCNE1* is significantly co-overexpressed with *TPX2* in (c) METABRIC: TNBC ($n = 299$, $p < 0.001$, $r = 0.65$) and in (d) TCGA: TNBC ($n = 115$, $p < 0.001$, $r = 0.63$)

Table 2 Clinical characteristics and outcomes of patients with *CCNE1*-amplified tumors in the “Discovery-TNBC” cohort

Patient ID	Age	Stage	(Neo)Adjuvant chemotherapy	Response to neoadjuvant chemotherapy	Adjuvant radiation	RFS (months)	OS (months)
1	50	I	Declined	N/A	No	58	N/A
2	50	III	TAC	ypT2N0Mx	No	11	23
3	46	III	AC Carbo/nab-paclitaxel	ypT3N1Mx	Yes	24	55
4	36	III	AC-T	ypT0N1Mx	Yes	13	18
5	35	III	Carbo/taxol	ypT3N2aMx	Yes	3	21
6	46	III	Carbo/taxol	ypT3N3aMx	Yes	13	43
7	50	II	Docetaxel, cisplatin	ypT1N1Mx	Yes	8	15
8	46	III	Carbo/taxol, AC	ypT2N2aMx	Yes	11	20
9	36	II	Carbo/nab-paclitaxel	ypT1cN0Mx	Yes	8	24
10	55	III	AC-T (adjuvant)	pT4N2aMx	No	46	60
11	46	II	Carbo/taxol (adjuvant)	pT2N1Mx	No	21	34
12	53	III	AC-T	ypT1N1aMx	No	9	16
13	48	II	TC	ypT1cN1Mx	Yes	3	38

AC adriamycin/cyclophosphamide, AC-T adriamycin/cyclophosphamide followed by paclitaxel, Carbo/taxol carboplatin/paclitaxel, TAC docetaxel/adriamycin/cyclophosphamide, TC docetaxel /cyclophosphamide, RFS relapse-free survival (surgery to first relapse), OS overall survival (surgery to death)



in TNBC is mutually exclusive with *BRCA1/2* mutations in both METABRIC and TCGA breast cancer databases.

Discussion

Tumor genomic evolution is frequently observed due to selection pressure imposed by chemotherapy. We

hypothesize that studying paired primary-metastatic TNBCs at the genomic and transcriptomic levels may lead to a better understanding of the underlying mechanisms behind chemotherapy resistance. Herein, we report mRNA expression profiling and targeted DNA sequencing data from paired primary-metastatic

TNBCs in a “Pilot-TNBC” cohort and an independent, expanded “Discovery-TNBC” cohort. Despite largely concordant gene expression levels, 8 putative cancer driver genes were significantly upregulated in metastatic samples compared to their matched primary specimens. Furthermore, *CCNE1* amplification was associated with poor overall survival in the “Discovery-TNBC” cohort, highlighting *CCNE1* amplification as a possible mechanism of chemotherapy resistance.

TNBCs are extremely heterogeneous. In an effort to better understand and categorize TNBCs, several molecular classifiers were developed (i.e. Lehmann/Pietenpol [3] and Burstein [4]). Although these classifiers have improved our understanding of TNBC tumor biology and helped distill scientific analyses into outputs that are more easily interpretable, none have led to changes in clinical practice. Herein, we identified so-called “molecular phenotype shifts” between select primary and metastatic samples from our “Pilot-TNBC” cohort, but a universal shift in TNBC pheno-subtypes was not observed. The clinical application of these findings requires further investigation.

Previous studies have reported variable findings upon comparison of primary and metastatic tumors. Some revealed concordant expression patterns between primary breast cancers and their matched synchronous lymph nodes [16, 35–37]. Vecchi, et al. found that expression of a specific gene set could differentiate between primary tumors and synchronous metastases [38]. However, Weigelt, et al. studied seven cases of primary breast cancer and asynchronous distant metastases and showed that a 70-gene prognostic signature was generally maintained in the switch from primary to metastasis across most of the pairs [39]. Several studies used targeted NGS to address genomic concordance between primary and metastases. Meric-Bernstam and colleagues reported that 86.6% of the somatic mutations and 62.3% of the copy number variations were concordant between primary tumors and recurrences [7]. These studies suggest that it may be necessary to study circulating tumor DNA (ctDNA), which is thought to be more representative of the mutational spectrum of the entirety of a patient’s metastatic disease rather than a biopsy of a single metastatic lesion [40]. Furthermore, non-coding RNAs and epigenetic modifications could also play an important role in the metastatic process [16, 41, 42].

Cyclin E and its associated CDK2 are essential for cellular progression through the G1 phase of the cell cycle and initiation of DNA replication. *CCNE1* amplification induces chromosome instability through persistent DNA replication and centrosome duplication [43–45]. *CCNE1*, along with its catalytic subunit CDK2, plays a critical role in cell cycle regulation to assure precise control of DNA replication, chromosome segregation and the G1 to S-phase transition [11, 12]. *CCNE1* amplification is

mutually exclusive with *BRCA1/2* mutations and correlates with cyclin E1 protein expression in ovarian cancer [46–48]. *CCNE1* amplification occurs in approximately 20% of high-grade serous ovarian carcinomas (HGSCs) [13, 14] and is associated with primary treatment resistance and poor outcome in these tumors [49]. Cyclin E1-specific pharmacological inhibitors are not yet available, although indirect *CCNE1* targeting trials are ongoing. However, small molecule inhibitors do exist for *CCNE1*’s counterpart in the cell cycle, CDK2. The pan CDK inhibitor, dinaciclib (SCH-727965), inhibits CDKs 2/5/1/9 and is being tested in clinical trials for hematological and solid malignancies (NCT00798213 and NCT00937937) [50, 51]. In addition, a WEE1 inhibitor AZD1755 is being tested in solid tumors with *CCNE1* amplification (NCT03253679). In a study of HGSC, expression of *TPX2*, a centromeric protein required for mitotic spindle function, was highly associated with *CCNE1* amplification [47]. In our study, *CCNE1* and *TPX2* were co-overexpressed in metastatic TNBCs compared to paired primaries. This finding is similar to previous results in ovarian cancer, where co-overexpression of *CCNE1* and *TPX2* were found to be related to clonal resistance to chemotherapy [47]. Mutual exclusivity of *BRCA1/2* mutation/deletion and *CCNE1* amplification has been shown previously in ovarian cancers [47]. Despite limited sample size, our finding of mutual exclusivity of *CCNE1* and *BRCA1/2* genes is worth further investigation and may help clinicians select patients for personalized treatment.

High cyclin E1 expression has been considered in other studies as a biomarker of poor clinical outcome in breast cancer [15, 52]. In a study of patients with HER2+ disease, *CCNE1* amplification/overexpression resulted in worse survival and CDK2 inhibition could overcome trastuzumab resistance in xenograft models [15]. Hunt, et al. demonstrated that overexpression of cytoplasmic low molecular weight cyclin E (LMW-E) is associated with poor survival in patients with breast cancer [53]. Because it lacks a nuclear localization domain, LMW-E accumulates in the cytoplasm, where it binds to CDK2 and retains kinase activity [54]. Our study provides further evidence of the association of *CCNE1* amplification and poor OS in patients with recurrent TNBCs. *CCNE1* mRNA was significantly upregulated in metastatic compared to paired primary TNBCs. Although no survival differences were observed, this may be reflective of the limited sample size.

This study features a combination of well-annotated clinicopathologic and genomic data, which allowed us to directly identify clinically relevant genomic information. However, our study is limited by sample size, lack of unified treatment variables, and retrospective nature. We did not study noncoding RNAs nor epigenetic changes within the tumors. Finally, although patient-matched, singular biopsies

do not represent a complete picture of metastatic disease. Overall, our study is hypothesis-generating and these results will benefit from verification with a larger cohort of patients and more comprehensive technologies and tumor biology assessments such as ctDNA analyses.

Conclusions

In this study, comparison of paired primary and metastatic TNBCs demonstrates heterogeneity of molecular mechanisms of gene expression underlying TNBC recurrence. Amplification of *CCNE1* is associated with poor OS in patients with metastatic TNBC. *CCNE1* amplification may serve as a target for therapeutic intervention in chemotherapy-resistant TNBCs.

Additional files

Additional file 1: Table S3. Genomic alteration data for the “Pilot-TNBC” cohort. (XLSX 39 kb)

Additional file 2: Figure S2. *CCNE1* is significantly co-overexpressed with *TPX2* in “Pilot-TNBC” and “Discovery-TNBC” cohorts ($p < 0.001$). (TIF 165 kb)

Additional file 3: Figure S1. TNBCs exhibited higher mRNA expression compared with non-TNBCs for the eight putative cancer driver genes identified in the differential expression analysis in (a) METABRIC (TNBC, $n = 299$ and non-TNBC, $n = 1605$) and (b) TCGA (TNBC, $n = 115$ and non-TNBC, $n = 976$). Analysis-values were calculated using the Wilcoxon test ($*p < 0.01$, $**p < 0.0001$); (c) TNBCs also exhibited higher mRNA expression compared to non-TNBCs and normal breast tissue for the eight putative cancer driver genes in TCGA (TNBC, $n = 115$; non-TNBC, $n = 976$; and normal breast, $n = 112$) as per the ANOVA test ($*p < 0.01$, $**p < 0.0001$), y-axis is the log2 of TPM values by RSEM. (TIF 120 kb)

Additional file 4: Table S1. Normalized mRNA expression data for the “Pilot-TNBC” cohort. (XLSX 13 kb)

Additional file 5: Table S2. Normalized mRNA expression data for the “Discovery-TNBC” cohort. (XLSX 20 kb)

Abbreviations

AC: Adriamycin/cyclophosphamide; AC-T: Adriamycin/cyclophosphamide followed by paclitaxel; BC: Breast cancer; BL1: Basal-like 1; BL2: Basal-like 2; Carbo/paclitaxel: Carboplatin/paclitaxel; *CCNE1*: Cyclin E1; cfDNA: Cell-free DNA; DEG: Differentially expressed genes; ER: Estrogen receptor; Exome-Seq: Exome-sequencing; FDR: False discovery rate; HER2+: Human epidermal growth factor receptor 2 positive; HGSCs: High-grade serous ovarian carcinomas; HQ: Highest quartile; HR+: Hormone-receptor positive; IDC: Invasive ductal carcinoma; ILC: Invasive lobular carcinoma; K-M: Kaplan-Meier; LAR: Luminal androgen receptor; LN: Lymph node; LQ: Lowest quartile; M: Mesenchymal; met: Metastatic tumor; METABRIC: Molecular Taxonomy of Breast Cancer International Consortium; mRNA: Messenger RNA; OS: Overall survival; PR: Progesterone receptor; RFS: Relapse-free survival; RMA: Robust multi-array average; TAC: Docetaxel/adriamycin/cyclophosphamide; TC: Docetaxel (Taxotere) /cyclophosphamide; TCGA: The Cancer Genome Atlas; TNBC: Triple negative breast cancer; VUS: Variants of unknown significance

Acknowledgements

This study was supported by STOP Cancer Foundation, NIH K-12 Career Development Award K12CA001727 (Joanne Mortimer MD), and National Cancer Institute P30CA034196, R21CA191848, and U24 CA224067 (Jeffrey H. Chuang PhD). Research reported in this publication included work performed in the Pathology and Biostatistics Core supported by the National Cancer Institute P30CA033572 Award. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The authors would thank Dr. Edison T. Liu and Dr. Francesca Menghi from The Jackson Laboratory for the discussion of data interpretation.

Funding

This study was supported by STOP Cancer Foundation, NIH K-12 Career Development Award K12CA001727 (Joanne Mortimer MD), and National Cancer Institute P30CA034196, R21CA191848, and U24 CA224067 (Jeffrey H. Chuang PhD).

Availability of data and materials

METABRIC and TCGA are publicly available databases. Normalized mRNA expression data for “Pilot-TNBC” cohort is available in Additional file 4: Table S1, normalized mRNA expression data for the “Discovery-TNBC” cohort is available in Additional file 5: Table S2, and genomic alteration data for the “Pilot-TNBC” cohort is available in Additional file 1: Table S3.

Authors’ contributions

ZMZ contributed to study conception, data interpretation, and manuscript revision; SEY contributed to data collection, data interpretation, and manuscript revision; KEH contributed to data interpretation and manuscript revision; SML contributed to data interpretation and manuscript revision; YCY contributed to data collection, data interpretation, and manuscript revision; JN contributed to data interpretation; ZL contributed to data interpretation; CW contributed to data interpretation; XW contributed to data interpretation; RMJ contributed to data analysis; JHC contributed to study conception, data interpretation, and manuscript revision; and YY contributed to study conception, data collection, data interpretation, and manuscript revision. All authors read and approved the final manuscript.

Authors’ information

Not applicable

Ethics approval and consent to participate

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all participants included in the study. All tumor specimens were identified through a City of Hope IRB-approved retrospective protocol from patients consented to City of Hope Biorepository Protocol.

Consent for publication

Not applicable

Competing interests

YY has contracted clinical trials and research projects sponsored by Merck, Eisai, Novartis, Genentech, and Pfizer. The other authors declare that they have no competing interests.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹The Jackson Laboratory for Genomic Medicine, Farmington, CT, USA. ²City of Hope Comprehensive Cancer Center and Beckman Research Institute, 1500 E. Duarte Road, Duarte, CA 91010, USA. ³Genentech, Inc., Oncology Biomarker Development, South San Francisco, CA, USA.

Received: 17 September 2018 Accepted: 7 January 2019

Published online: 21 January 2019

References

- Collignon J, Lousberg L, Schroeder H, Jerusalem G. Triple-negative breast cancer: treatment challenges and solutions. *Breast Cancer*. 2016;8:93.
- Zeichner SB, Terawaki H, Gogineni K. A review of systemic treatment in metastatic triple-negative breast cancer. *Breast Cancer*. 2016;10:BCBCR. S32783.
- Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, Pietenpol JA. Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest*. 2011;121(7):2750–67.

4. Burstein MD, Tsimelzon A, Poage GM, Covington KR, Contreras A, Fuqua SA, Savage MI, Osborne CK, Hilsenbeck SG, Chang JC. Comprehensive genomic analysis identifies novel subtypes and targets of triple-negative breast cancer. *Clin Cancer Res*. 2014.
5. Lehmann BD, Jovanovic B, Chen X, Estrada MV, Johnson KN, Shyr Y, Moses HL, Sanders ME, Pietersen JA. Refinement of triple-negative breast cancer molecular subtypes: implications for neoadjuvant chemotherapy selection. *PLoS One*. 2016;11(6):e0157368.
6. Gao R, Davis A, McDonald TO, Sei E, Shi X, Wang Y, Tsai P-C, Casasent A, Waters J, Zhang H. Punctuated copy number evolution and clonal stasis in triple-negative breast cancer. *Nat Genet*. 2016;48(10):1119.
7. Meric-Bernstam F, Frampton GM, Ferrer-Lozano J, Yelensky R, Pérez-Fidalgo JA, Wang Y, Palmer GA, Ross JS, Miller VA, Su X. Concordance of genomic alterations between primary and recurrent breast cancer. *Mol Cancer Ther*. 2014;13(5):1382–9.
8. Liedtke C, Brogi K, Moulder S, Hsu L, Kau S-W, Symmans W, Albarracín C, Meric-Bernstam F, Woodward W, Theriault R. Prognostic impact of discordance between triple-receptor measurements in primary and recurrent breast cancer. *Ann Oncol*. 2009;20(12):1953–8.
9. Niikura N, Liu J, Hayashi N, Mittendorf EA, Gong Y, Palla SL, Tokuda Y, Gonzalez-Angulo AM, Hortobagyi GN, Ueno NT. Loss of human epidermal growth factor receptor 2 (HER2) expression in metastatic sites of HER2-overexpressing primary breast tumors. *J Clin Oncol*. 2011;30(6):593–9.
10. Jensen JD, Laenkholm A-V, Knoop A, Ewertz M, Bandaru R, Liu W, Hackl W, Barrett JC, Gardner H. PIK3CA mutations may be discordant between primary and corresponding metastatic disease in breast cancer. *Clin Cancer Res*. 2011;17(4):667–77.
11. Sapoznik S, Aviel-Ronen S, Bahar-Shany K, Zadok O and Levanon K. CCNE1 expression in high grade serous carcinoma does not correlate with chemoresistance. *Oncotarget* 2017; 8(37):62240.
12. Bendris N, Lemmers B, Blanchard JM. Cell cycle, cytoskeleton dynamics and beyond: the many functions of cyclins and CDK inhibitors. *Cell Cycle*. 2015; 14(12):1786–98.
13. Karst AM, Jones PM, Vena N, Ligon AH, Liu JF, Hirsch MS, Etemadmoghadam D, Bowtell DD, Drapkin R. Cyclin E1 deregulation occurs early in secretory cell transformation to promote formation of fallopian tube-derived high-grade serous ovarian cancers. *Cancer Res*. 2014;74(4): 1141–52.
14. Kuhn E, Wang TL, Doberstein K, Bahadiri-Talbot A, Ayhan A, Sehdev AS, Drapkin R, Kurman RJ, Shih le M. CCNE1 amplification and centrosome number abnormality in serous tubal intraepithelial carcinoma: further evidence supporting its role as a precursor of ovarian high-grade serous carcinoma. *Mod Pathol*. 2016;29(10):1254–61.
15. Scaltriti M, Eichhorn PJ, Cortés J, Prudkin L, Aura C, Jiménez J, Chandralapaty S, Serra V, Prat A, Ibrahim YH. Cyclin E amplification/overexpression is a mechanism of trastuzumab resistance in HER2+ breast cancer patients. *Proc Natl Acad Sci*. 2011;108(9):3761–6.
16. Krøigård AB, Larsen MJ, Thomassen M, Kruse TA. Molecular concordance between primary breast cancer and matched metastases. *Breast J*. 2016; 22(4):420–30.
17. Yates LR, Gerstung M, Knappskog S, Desmedt C, Gundem G, Van Loo P, Aas T, Alexandrov LB, Larsimont D, Davies H. Subclonal diversification of primary breast cancer revealed by multiregion sequencing. *Nat Med*. 2015;21(7):751.
18. Network CGA. Comprehensive molecular portraits of human breast tumours. *Nature*. 2012;490(7418):61.
19. Curtis C, Shah SP, Chin S-F, Turashvili G, Rueda OM, Dunning MJ, Speed D, Lynch AG, Samarajiwa S, Yuan Y. The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature*. 2012; 486(7403):346.
20. Bolstad BM, Irizarry RA, Astrand M, Speed TP. A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics (Oxford, England)*. 2003;19(2):185–93.
21. Von Minckwitz G, Hahnen E, Fasching PA, Hauke J, Schneeweiss A, Salat C, Reza M, Blohmer JU, Zahm DM and Jackisch C. Pathological complete response (pCR) rates after carboplatin-containing neoadjuvant chemotherapy in patients with germline BRCA (g BRCA) mutation and triple-negative breast cancer (TNBC): Results from GeparSixto. 2014;1005–1005. http://ascopubs.org/doi/abs/10.1200/jco.2014.32.15_suppl.1005.
22. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res*. 2015;43(7):e47–e47.
23. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B Methodol*. 1995;289–300.
24. Tutt A, Ellis P, Kilburn L, Gilett C, Pinder S, Abraham J, Barrett S, Barrett-Lee P, Chan S, Cheang M, Dowsett M, Fox L, Gazinska P, Grigoriadis A, Gutin A, Harper-Wynne C, et al. Abstract S3-01: the TNT trial: a randomized phase III trial of carboplatin (C) compared with docetaxel (D) for patients with metastatic or recurrent locally advanced triple negative or BRCA1/2 breast cancer (CRUK/07/012). *Cancer Res*. 2015;75(9 Supplement):S3-01-S03-01.
25. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2010;26(1):139–40.
26. McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res*. 2012;40(10):4288–97.
27. Frampton GM, Fichtenholtz A, Otto GA, Wang K, Downing SR, He J, Schnall-Levin M, White J, Sanford EM, An P. Development and validation of a clinical cancer genomic profiling test based on massively parallel DNA sequencing. *Nat Biotechnol*. 2013;31(11):1023.
28. Ross JS, Badve S, Wang K, Sheehan CE, Boguniewicz AB, Otto GA, Yelensky R, Lipson D, Ali S, Morosini D. Genomic profiling of advanced-stage, metaplastic breast carcinoma by next-generation sequencing reveals frequent, targetable genomic abnormalities and potential new treatment options. *Arch Pathol Lab Med*. 2015;139(5):642–9.
29. Balko JM, Giltman J, Wang K, Schwarz LJ, Young CD, Cook RS, Owens P, Sanders ME, Kuba MG, Sánchez V. Molecular profiling of the residual disease of triple-negative breast cancers after neoadjuvant chemotherapy identifies actionable therapeutic targets. *Cancer Discov*. 2013;CD-13-0286.
30. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW. Cancer genome landscapes. *Science*. 2013;339(6127):1546–58.
31. Lawrence MS, Stojanov P, Mermel CH, Robinson JT, Garraway LA, Golub TR, Meyerson M, Gabriel SB, Lander ES, Getz G. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature*. 2014;505(7484): 495.
32. Zhao Z-M, Zhao B, Bai Y, Iamarino A, Gaffney SG, Schlessinger J, Lifton RP, Rimm DL, Townsend JP. Early and multiple origins of metastatic lineages within primary tumors. *Proc Natl Acad Sci*. 2016;113(8):2140–5.
33. Griffith M, Griffith OL, Coffman AC, Weible JV, McMichael JF, Spies NC, Koval J, Das I, Callaway MB, Eldred JM. DGI: mining the druggable genome. *Nat Methods*. 2013;10(12):1209.
34. Van Allen EM, Wagle N, Stojanov P, Perrin DL, Cibulskis K, Marlow S, JaneValbuena J, Friedrich DC, Kryukov G, Carter SL. Whole-exome sequencing and clinical interpretation of formalin-fixed, paraffin-embedded tumor samples to guide precision cancer medicine. *Nat Med*. 2014;20(6):682.
35. Ellsworth RE, Seebach J, Field LA, Heckman C, Kane J, Hooke JA, Love B, Shriver CD. A gene expression signature that defines breast cancer metastases. *Clin Exp Metastasis*. 2009;26(3):205–13.
36. Suzuki M, Tarin D. Gene expression profiling of human lymph node metastases and matched primary breast carcinomas: clinical implications. *Mol Oncol*. 2007;1(2):172–80.
37. Hao X, Sun B, Hu L, Lähdesmäki H, Dunmire V, Feng Y, Zhang SW, Wang H, Wu C, Wang H. Differential gene and protein expression in primary breast malignancies and their lymph node metastases as revealed by combined cDNA microarray and tissue microarray analysis. *Cancer*. 2004; 100(6):1110–22.
38. Vecchi M, Confalonieri S, Nuciforo P, Viganò M, Capra M, Bianchi M, Nicosia D, Bianchi F, Galimberti V, Viale G. Breast cancer metastases are molecularly distinct from their primary tumors. *Oncogene*. 2008;27(15):2148.
39. Weigelt B, Hu Z, He X, Livasy C, Carey LA, Ewend MG, Glas AM, Perou CM, Van't Veer LJ. Molecular portraits and 70-gene prognosis signature are preserved throughout the metastatic process of breast cancer. *Cancer Res*. 2005;65(20):9155–8.
40. Rothe F, Laes J-F, Lambrechts D, Smeets D, Vincent D, Maetens M, Fumagalli D, Michiels S, Dris S, Moerman C. Plasma circulating tumor DNA as an alternative to metastatic biopsies for mutational analysis in breast cancer. *Ann Oncol*. 2014;25(10):1959–65.
41. Gravgaard KH, Lyng MB, Laenkholm A-V, Søkilde R, Nielsen BS, Litman T, Ditzel HJ. The miRNA-200 family and miRNA-9 exhibit differential expression in primary versus corresponding metastatic tissue in breast cancer. *Breast Cancer Res Treat*. 2012;134(1):207–17.

42. Chisholm KM, Wan Y, Li R, Montgomery KD, Chang HY, West RB. Detection of long non-coding RNA in archival tissue: correlation with polycomb protein expression in primary and metastatic breast carcinoma. *PLoS One*. 2012;7(10):e47998.
43. Ekholm-Reed S, Méndez J, Tedesco D, Zetterberg A, Stillman B, Reed SI. Deregulation of cyclin E in human cells interferes with prereplication complex assembly. *J Cell Biol*. 2004;165(6):789–800.
44. Kawamura K, Izumi H, Ma Z, Ikeda R, Moriyama M, Tanaka T, Nojima T, Levin LS, Fujikawa-Yamamoto K, Suzuki K. Induction of centrosome amplification and chromosome instability in human bladder cancer cells by p53 mutation and cyclin E overexpression. *Cancer Res*. 2004;64(14):4800–9.
45. Nakayama N, Nakayama K, Shamima Y, Ishikawa M, Katagiri A, Iida K, Miyazaki K. Gene amplification CCNE1 is related to poor survival and potential therapeutic target in ovarian cancer. *Cancer: interdisciplinary international journal of the. Am Cancer Soc*. 2010;116(11):2621–34.
46. Yang L, Fang D, Chen H, Lu Y, Dong Z, Ding H-F, Jing Q, Su S-B, Huang S. Cyclin-dependent kinase 2 is an ideal target for ovary tumors with elevated cyclin E1 expression. *Oncotarget*. 2015;6(25):20801.
47. Etemadmoghadam D, Weir B, Au-Yeung G, Alsop K, Mitchell G and George J. Australian ovarian Cancer study group Davis S, D'Andrea AD, Simpson K, Hahn WC, Bowtell DD. Synthetic lethality between CCNE1 amplification and loss of BRCA1. *Proc Natl Acad Sci US A* 2013; 110:19489–19494.
48. Nakayama N, Nakayama K, Shamima Y, Ishikawa M, Katagiri A, Iida K, Miyazaki K. Gene amplification CCNE1 is related to poor survival and potential therapeutic target in ovarian cancer. *Cancer*. 2010;116(11):2621–34.
49. Au-Yeung G, Lang F, Azar WJ, Mitchell C, Jarman KE, Lackovic K, Aziz D, Cullinane C, Pearson RB, Mileskin L. Selective targeting of cyclin E1-amplified high-grade serous ovarian cancer by cyclin-dependent kinase 2 and AKT inhibition. *Clin Cancer Res*. 2017;23(7):1862–74.
50. Mitri Z, Karakas C, Wei C, Briones B, Simmons H, Ibrahim N, Alvarez R, Murray JL, Keyomarsi K, Moulder S. A phase 1 study with dose expansion of the CDK inhibitor dinaciclib (SCH 727965) in combination with epirubicin in patients with metastatic triple negative breast cancer. *Investig New Drugs*. 2015;33(4):890–4.
51. Kumar SK, LaPlant B, Chng WJ, Zonder J, Callander N, Fonseca R, Fruth B, Roy V, Erlichman C, Stewart AK. Dinaciclib, a novel CDK inhibitor, demonstrates encouraging single-agent activity in patients with relapsed multiple myeloma. *Blood*. 2015;125(3):443–8.
52. Keyomarsi K, Tucker SL, Buchholz TA, Callister M, Ding Y, Hortobagyi GN, Bedrosian I, Knickerbocker C, Toyofuku W, Lowe M. Cyclin E and survival in patients with breast cancer. *N Engl J Med*. 2002;347(20):1566–75.
53. Hunt KK, Karakas C, Ha MJ, Biernacka A, Yi M, Sahin AA, Adjapong O, Hortobagyi GN, Bondy ML, Thompson PA, Cheung KL. Cytoplasmic cyclin E predicts recurrence in patients with breast cancer *Clin Cancer Res*. 2017; 23(12):2991–3002. <http://clincancerres.aacrjournals.org/content/23/12/2991>.
54. Delk NA, Hunt KK, Keyomarsi K. Altered subcellular localization of tumor-specific cyclin E isoforms affects cyclin-dependent kinase 2 complex formation and proteasomal regulation. *Cancer Res*. 2009;69(7):2817–25.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

